



Presentation

100% w/w/ oxytetracycline premix for top dressing of already manufactured feeding stuffs.

Dosage and Administration

Aquatet is given by oral administration at the rate of 75 mg/kg bodyweight of fish daily for initially 4 days. After the initial treatment, mortalities should be assessed and a further 4 day treatment course should be administered if necessary. Aquatet is for administration only through the feed by mixing with already manufactured feed. Feeding rates will vary according to water temperature and it may therefore be more convenient to medicate on a basis of a fixed rate e.g. 1% of bodyweight, with an extra daily feed requirement being met by unmedicated food. The following inclusion rates will provide the recommended dose:

% bodyweight	Daily feed rate Aquatet inclusion rate		
	per 5 kg of food	per 25 kg of food	per 1 tonne of food
0.5	75 g	375 g	15 kg
1	37.5 g	187.5 g	7.5 kg
2	18.75 g	93.75 g	3.75 kg

- 100% w/w active oxytetracycline
- Effective against bacteria: Aeromonas spp, Yersinia spp, Vibrio spp and Myxobacteria spp
- For use in Atlantic Salmon and Rainbow trout
- 100% w/w premix for food
- 720° days withdrawal time

Method of mixing:

Weigh out appropriate amounts of fish pellets and Aquatet powder and mix well together in the dry state. As an aid to the adhesion of the Aquatet powder in the fish pellets, a small quantity of tepid gelatin solution or edible oil is then added to the food while mixing. Sufficient quantity should be mixed into the medicated food until it is slightly dampened.

Legal Category

POM-V

Package Quantities

1 kg, 2.5 kg plastic tubs

Marketing Authorisation Number

Vm 11003/4002

Marketing Authorisation Holder

PHARMAQ Ltd
Unit 15, Sandheath Industrial Estate,
Fordingbridge, Hampshire SP6 1PA
Tel No: 01425 656081
Fax: 01425 657992
Website: www.pharmaq.no

Contra-indications, Warnings etc.

Animals must not be slaughtered for human consumption during treatment. Fish may be slaughtered for human consumption 720° days after the end of treatment.

Avoid skin contact. Whilst handling the product wear coveralls, protective goggles and chemically resistant impermeable gloves at all times. Avoid inhalation of dust. Wear either a disposable half-mask respirator conforming to European Standard EN 149, or a non-disposable respirator European Standard EN 140 with a filter to EN 143.

Pharmaceutical Precautions

Store the original container at a temperature not exceeding 25°C and in a dry place. Re-close the part-used containers. Protect from light. Medicated diet should be used for the treatment period only. Keep out of the reach and sight of children.

Licensed in

United Kingdom and Ireland.
Only available through your veterinarian.

